



**GOAT ANTI-MOUSE IgM+IgG+IgA(H+L)  
HRP CONJUGATED, SPECIES ADSORBED: HUMAN  
POLYCLONAL ANTIBODY**

<b>CATALOG NUMBER:</b>	AP501P	<b>QUANTITY:</b>	1.0 mL
<b>LOT NUMBER:</b>			
<b>EPITOPE:</b>	Mouse IgM, IgG, IgA	<b>HOST/ISOTYPE:</b>	Goat
<b>BACKGROUND:</b>	IgG is the most abundant immunoglobulin composed of two heavy chains and two light chains. Each molecule has two antigen binding sites. IgG has 4 subclasses: IgG1 (66%), IgG2 (23%), IgG3 (7%) and IgG4 (4%). IgM constitutes about 10% of serum immunoglobulins. IgM (with IgD) is the major immunoglobulin expressed on the surface of B cells. Monomeric IgA constitutes 5-15 % of the serum immunoglobulins whereas dimeric IgA is localized to mucosa surfaces.		
<b>SPECIFICITY:</b>	The antibody reacts with the heavy and light chains of mouse IgM, IgG1, IgG2a, IgG2b, IgG3 and IgA as demonstrated by ELISA and flow cytometry. Minimal cross reactivity with human immunoglobulins.		
<b>APPLICATIONS:</b>	<u>ELISA:</u> 1:4,000-1:8,000 <i>Optimal working dilutions must be determined by the end user.</i>		
<b>SPECIES REACTIVITY:</b>	Reacts with heavy and light chains of mouse IgM, IgG1, IgG2a, IgG2b, IgG3 and IgA. Absorbed for human sera and purified human paraproteins. Minimal cross reactivity with human immunoglobulins. Reactivity with other species has not been determined.		
<b>IMMUNOGEN:</b>	Prepared from pooled antisera from goats hyperimmunized with mouse IgM, IgG and IgA paraproteins.		
<b>PRESENTATION:</b>	Purified by affinity chromatography on pooled mouse IgM+IgG+IgA covalently linked to agarose. Liquid in 1.0 mL of stock solution in 50% glycerol/50% PBS, pH 7.4.		
<b>STORAGE/HANDLING:</b>	Store long-term at -20°C. Maintain refrigerated at 2°-8°C under sterile conditions for up to twelve months from date of receipt.		

**Important Note:** *During shipment, small volumes of product will occasionally become entrapped in the seal of the product vial. For products with volumes of 200 µL or less, we recommend gently tapping the vial on a hard surface or briefly centrifuging the vial in a tabletop centrifuge to dislodge any liquid in the container's cap.*

**FOR RESEARCH USE ONLY; NOT FOR USE IN DIAGNOSTIC  
PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION**

Unless otherwise stated in our catalog or other company documentation accompanying the product(s), our products are intended for research use only and are not to be used for any other purpose, which includes but is not limited to, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses or any type of consumption or application to humans or animals.

©2002 - 2011: Millipore Corporation. All rights reserved. No part of these works may be reproduced in any form without permission in writing.